

SURVEILLANCE OF INFECTIOUS DISEASES IN ANIMALS AND HUMANS IN SWEDEN 2019

Chapter excerpt -
Infectious diseases in fish, crustaceans and molluscs



Editor: Karl Ståhl

Department of Disease Control and Epidemiology, National Veterinary Institute (SVA), SE-751 89 Uppsala, Sweden.

Authors: Charlotte Axén, Mia Brytting, Ioana Bujila, Erika Chenais, Rikard Dryselius, Helena Eriksson, Pernille Etterlin, Eva Forsgren, Malin Grant, Gittan Gröndahl, Gunilla Hallgren, Anette Hansen, Gete Hestvik, Marika Hjertqvist, Mia Holmberg, Cecilia Hultén, Helena Höök, Cecilia Jernberg, Jerker Jonsson, Ulrika König, Elina Lahti, Emelie Larsdotter, Mats Lindblad, Anna Lundén, Emma Löf, Hans-Olof Nilsson, Oskar Nilsson, Maria Nöremark, Anna Ohlson, Ylva Persson, Karin Persson-Waller, Thomas Rosendal, Karl Ståhl, Robert Söderlund, Kaisa Sörén, Magnus Thelander, Karin Troell, Anders Wallensten, Per Wallgren, Stefan Widgren, Ulrika Windahl, Beth Young, Nabil Yousef, Siamak Zohari, Erik Ågren, Estelle Ågren, Elina Åsbjer

Cover Photo: Anders Lindström

Copyright of map data: ©EuroGeographics for the administrative boundaries

Reporting guidelines: Reporting guidelines were introduced in 2018 for those those chapters related to purely animal pathogens. The guidelines build on experiences from several EU projects, and have been validated by a team of international experts in animal health surveillance. The aim is to develop these guidelines further in collaboration within the global surveillance community and they have therefore been made available in the form of a wiki on the collaborative platform GitHub (<https://github.com/SVA-SE/AHSURED/wiki>). Feel free to contribute!

Layout: The production of this report continues to be accomplished using a primarily open-source toolset. The method allows the source text, produced by authors, to be edited independently of the template for the layout which can be modified and reused for future reports. Specifically, the chapter texts, tables and captions are authored in Microsoft Word and then converted using pandoc and R to the LaTeX typesetting language. Most figures and maps are produced using the R software for statistical computing. Development for 2019 has further improved the importing of content from Word to LaTeX. The method can now import text, tables and figure captions from Word, as well as the newly designed 'IN FOCUS' sections of some chapters. The tool is available as an R-package at GitHub (<https://github.com/SVA-SE/mill/>). This year the report was also built with a continuous integration pipeline on Microsoft's Azure DevOps platform, allowing every committed change to the content to be built and tested automatically. The report generation R-package and process was designed by Thomas Rosendal and Stefan Widgren. In 2019, figures and the final typesetting were done by Wiktor Gustafsson and Thomas Rosendal with contributions from the report authors.

Print: TMG Tabergs AB.

Text, tables, figures and maps may be cited and reprinted only with reference to this report.

Suggestion citation: Surveillance of infectious diseases in animals and humans in Sweden 2019, National Veterinary Institute (SVA), Uppsala, Sweden. SVA:s rapportserie 64 1654-7098.

This report may be subject to updates and corrections. The latest version is always available for download at www.sva.se.

Infectious diseases in fish, crustaceans and molluscs



Juvenile rainbow trout with suspicion of flavobacteriosis. Photo: SVA.

BACKGROUND

All registered aquaculture farming sites are obligated to participate in the Official Health Control Programme, regulated in accordance with SJVFS 2014:4 and by Council Directive 2006/88/EG. Sweden has a very good health status in aquaculture as well as in wild populations of fish and shellfish. None of the serious diseases that occur throughout Europe are prevalent in Sweden. A restrictive approach to imports of live fish for restocking/farming, an early introduction of health control in farms and the presence of hydroelectric dams in most Swedish rivers (acting as migration barriers for feral fish from the coastal zone) all contribute to maintaining this health status. The presence of dams also results in a different health status at the coast compared to the more disease-free continental zone. To maintain this situation, all transport of live fish from the coast to the inland zone is forbidden and Sweden has a national restocking programme for salmonids to compensate for the lack of natural migration.

LEGISLATION AND DISEASES

All Swedish fish farms have participated in surveillance for the diseases mentioned below since the late 1980s in accordance with Commission Decision 2001/183/EC (now replaced by Commission Implementing Decision (EU) 2015/1554) and Council Directive 2006/88/EC. Sweden has an approved disease-free zone status (2002/308/EC) for Viral haemorrhagic septicaemia (VHS) and Infectious haematopoietic necrosis (IHN) (2008/427/EC). Additional

guarantees are in place for the whole country for Spring Viraemia of Carp (SVC), and for the inland zone for Infectious Pancreatic Necrosis (IPN) (2010/221/EU). The inland zone has an eradication programme for Renibacteriosis/bacterial kidney disease (BKD) and the coastal zone for IPN (2010/221/EU). These diseases are included in the Swedish legislation on notifiable diseases (SJVFS 2013:23). Further, IHN, VHS, IPN (other than serotype ab) and SVC are included in the Swedish Act on epizootic diseases (SFS 1999:657 with amendments). In addition, testing is routinely done for Koi herpes virus (KHV) in imported, quarantined koi, and for Crayfish plague in crayfish. These diseases are also regulated by the Swedish legislation on notifiable diseases (SJVFS 2013:23). Other notifiable diseases such as furunculosis (*Aeromonas salmonicida salmonicida*/ASS), yersiniosis/Enteric redmouth disease (ERM), Marteiosis and Bonamiosis (shellfish) and Whitespot disease (crayfish) are not actively tested for within surveillance programmes.

Infectious haematopoietic necrosis (IHN) and viral haemorrhagic septicaemia (VHS)

Both diseases are caused by rhabdoviruses and occur frequently in Europe. They are transferred horizontally, but vertical transmission cannot be completely ruled out for IHN. Both diseases have greatest impact in freshwater rainbow trout (*Oncorhynchus mykiss*) aquaculture but have also been detected in several other species. Infected fish exhibit behavioural changes, lethargy and abnormal swimming

(whirling). The fish are anaemic with varying degrees of haemorrhage in multiple organs. VHS also exists in a marine form, and a low prevalence in wild populations of sensitive species cannot be excluded in the Swedish coastal zone since the virus has been identified in wild fish from Skagerrak and the Bornholm basin. IHN was found in two Bothnian bay farms in Finland in 2017, but the virus has not yet been identified in Sweden.

Infectious pancreatic necrosis (IPN)

IPN is caused by a birnavirus which is highly infectious to juvenile salmonids. Susceptibility declines with increasing age. Fish that survive infection become subclinical carriers. In addition to salmonids, the virus has been detected in several other species. The virus is transmitted both horizontally and vertically.

There are seven genogroups with varying virulence. Some genogroups cause up to 90% mortality in fry, and IPN is considered one of the costliest fish diseases in several European countries. Symptoms include darkening, abdominal distension and corkscrew swimming. Petechial haemorrhage in abdominal fat and internal organs are the most common internal findings. IPN appears sporadically in Swedish east coast farms.

Renibacteriosis (BKD)

BKD is caused by a gram-positive bacterium, *Renibacterium salmoninarum*. The infection can be transmitted both horizontally and vertically. The disease favours low water temperatures, and outbreaks mainly occur at temperatures between 7 and 15°C.

Salmon and arctic char are most susceptible to BKD and mortality can reach 80%. In rainbow trout, the disease is chronic with a continuous low mortality of about 5–10%, however outbreaks of up to 40% mortality can occur. Infected fish may have reduced growth and disease can result in a deterioration of the meat quality. BKD is present in a few farms in the Swedish inland zone.

Spring viraemia of carp (SVC)

SVC is caused by a rhabdovirus. The disease occurs in Asia and several European countries. SVC is not present in Sweden. The virus has been detected in several fish species in the cyprinid family and is transmitted horizontally. Clinical signs are usually general, such as darkening, exophthalmia and a slow breathing. The fish swim lazily with sporadic periods of hyperactivity. Other common findings are pale gills, ascites and haemorrhages in the skin and gills. Internally, haemorrhage is found in various organs including muscle, swim bladder and the brain.

Koi herpes virus (KHV) infection

KHV is a herpesvirus and affects common carp (*Cyprinus carpio*) and variants thereof, including koi. The virus was first detected in 1998 and has since then been reported from all continents except Australia. The virus is transmitted horizontally. KHV can cause severe problems and is associated with high mortality. Infected fish usually swim at the surface and have an increased breathing frequency. Symptoms include enophthalmia, gill necrosis and secondary bacterial or parasitic infections on gills and skin. Surviving carps can

become subclinical carriers. The prevalence in Sweden is unknown. Koi is frequently imported, but only a few farms use quarantine and sampling. There were two Swedish outbreaks with 90–100% mortality in 2018.

Crayfish plague

Crayfish plague is caused by an aquatic fungus (*Aphanomyces astaci*) that spread from the United States to Europe in the late 1800s with live crayfish. The disease occurs throughout Europe and North America. The fungus reproduces by spores spread in the water. When the spores infect crayfish, they grow through the skin and attack the underlying tissues.

The signal crayfish becomes subclinically infected and may exhibit black (melanised) areas in the shell adjacent to the presence of the fungus in the skin. The spots disappear when the shell is shed but may gradually reappear.

When noble crayfish are infected, the first sign is high mortality in affected populations. Disease in the individual is characterised by behavioural changes such as movement during daytime, reduced coordination and balance difficulties.

Crayfish plague is spread in the southern parts of Sweden.

White spot disease (WSD)

WSD is caused by a *Whispovirus* (WSSv) that can infect a wide range of aquatic crustaceans, including marine, brackish and freshwater prawns, crabs, crayfish and lobsters. Outbreaks occur at water temperatures of 18–30°C. The most common clinical sign is white spots in the exoskeleton, but the disease can occur without obvious external signs.

The virus is transmitted both horizontally and vertically and has a long survival time outside the host animal. The virus can be present in imported frozen raw giant shrimps. There is a non-negligible risk that the virus will be introduced to the aquatic environment by anglers using these shrimps for bait. The consequences are difficult to predict but the virus may have a negative impact on Swedish crustacean populations. WSSv has never been detected in Sweden.

Marteiliosis

Marteiliosis, a disease in oysters and blue mussels, is caused by a unicellular parasite (*Marteilia refringens* in oysters and *M. pararefringens* in blue mussels). The parasite needs a crustacean (*Paracartia grani*) as an intermediate host. The disease causes reduced fitness, impaired growth and resorption of the gonads and hence reduced reproductive capacity. *M. pararefringens* is present on the Swedish west coast.

Bonamiosis

Bonamiosis is a disease in oysters caused by the protistan parasite *Bonamia ostreae*. The parasite invades and destroys the haemocytes. Usually, the only sign of disease is increased mortality in the infected oyster population. *B. ostreae* is found along the European Atlantic coast as far up as Denmark, where it has now been found in Limfjorden.

SURVEILLANCE

Within the Official Health Control Programme, there is active surveillance for the viruses causing IHN, VHS, IPN and SVC, and for renibacteriosis/BKD. Sampling frequency is based on classification of each farm into one of three categories (high (I), medium (II) or low risk (III)) after a risk analysis, based on the risk for the farm becoming infected, the risk that the farm will further spread the pathogen and the impact of the pathogen. The risk categorisation is performed by the Swedish Board of Agriculture. Farms within risk category I and II are tested every year and every second year, respectively, but farms within risk category III are only tested upon suspicion of disease. The aim of the Official Health Control Programme is to document freedom from disease and to contribute to the maintenance of this status.

There is also active surveillance in imported quarantined fish (eel - IPN and koi/carp - KHV). Active surveillance is also done when potential invasive alien species - like the marble crayfish - are discovered.

Passive disease surveillance has been done through diagnostics related to disease outbreaks in farms and wild fish.

Crayfish plague is monitored by passive surveillance and testing is done based on suspicion of disease outbreaks.

DIAGNOSTIC PROCEDURES

All diagnostic virus analyses are performed according to recommendations by the EU (EU 2015/1554) or the OIE aquatic manual at the Swedish reference laboratory for fish diseases at the National Veterinary Institute. Pooled organ material (for VHS, IHN and IPN spleen, kidney, heart/brain are tested, for SVC spleen, kidney, brain and gill are tested) by a cell culturing method. A pool consists of organs from up to ten fish (up to five fish for SVC). A cell culture is defined as virus-positive if a cytopathogenic effect is detected within two weeks, after which the virus is identified by ELISA and confirmed by real-time PCR (rt-PCR), or in some cases by serum neutralisation (SN) test. Thirty fish are sampled in regular fish farms, and in restocking farms all females are sampled after stripping of roe. In eel quarantine, 120 glass eels are sampled at arrival, and after two months 120 co-habituated rainbow trout are sampled for detection of virus. In the case of carp/koi, only a few fish may be sampled. KHV is tested on individual fish (pooled gill and kidney) by PCR.

BKD is tested on kidney tissue from individual fish and demonstrated by an ELISA method. Verification is done by rt-PCR. Thirty fish are sampled in regular farms, and in restocking farms up to 120 fish (all females) are sampled after stripping of roe.

A. astaci is demonstrated by light microscopy and cultivation and verified by rt-PCR, and WSSv is detected by rt-PCR. The number of sampled animals varies from case to case.

RESULTS

Official health control programme for fish farms and crustacean surveillance

The number of samples analysed and results are shown in Table 33. In summary, the active surveillance detected six cases of BKD and one case of IPN (one case=one outbreak). Four of six BKD cases were reinfections of recently sanitised farms.

Voluntary health control programme for fish farms

There were five recorded outbreaks of “other” notifiable diseases in fish during 2019. Furunculosis (ASS) was detected in four farms. One farm had recurrent disease and concurrent BKD infection, and another farm within the same company also got infected. The other two farms were unrelated cases. Yersiniosis was detected in one sea based farm.

Few cases (n=6) of flavobacteriosis due to *Flavobacterium psychrophilum*, usually the predominant production disease, were detected compared to previous years (15–30 cases in the last 10 years). Instead, *Aeromonas* bacteria other than *A. salmonicida salmonicida* or *A. salmonicida* atypically dominated with 16 cases. The cause for this shift is unknown. *Flavobacterium columnare* was detected in five disease cases during the summer.

Voluntary health control in wild fish

Due to the detection of IPN genogroup 6 in broodstock trout from Lake Vänern in December 2016, the County Boards of Värmland and Västra Götaland took an initiative to sample 500 wild trout and salmon from the lake in 2017. During 2017–2018, a total of 609 fish caught in the lake or broodstock that returned from the lake to the river and five organ pools with unknown number of fish (n=1–10) were investigated by virus cell culture. No virus was detected. Since only 279 of the 609 fish were caught in the lake, one more sampling was performed in 2019. A total of 41 fish were investigated for presence of virus in 2019. No virus was detected.

Outbreaks in wild fish, crustaceans and molluscs

One arctic char that had been caught in lake Vättern 2018 (frozen after catch) and had multiple small skin wounds was suspected to have spawning rash, a specific form of BKD, was confirmed to be infected.

Suspicion of crayfish plague was investigated in five outbreaks of mortality in noble crayfish and three cases with signal crayfish. Crayfish plague was not detected in any of the noble crayfish cases. Crayfish plague was detected in two of the signal crayfish cases. One case concerned collapse of a signal crayfish population and it cannot be excluded that the plague was involved, although signal crayfish are generally resistant to clinical infection. One case was a follow up from a water system where the plague had previously been identified. In the third case, neither crayfish plague nor any other pathogen was detected. The crayfish was sent in because the meat had turned black at boiling.

One case of mortality in Norwegian lobsters was also handled. A few different bacteria were identified, and investigation of fixed shell with underlying tissue identified severe thinning of the cuticula and widespread muscular necrosis. No certain diagnosis could be made, but idiopathic muscle necrosis with secondary bacterial infection was suspected.

DISCUSSION

The number of farms that were sampled are listed in Table 33. Swedish aquaculture has a good health status, where all severe diseases of EU/OIE importance are absent.

The most problematic disease to control is renibacteriosis/BKD, due to its vertical transmission and variable clinical presentation. Prolonged time from diagnosis to slaughter can lead to secondary health issues and increased antibiotic use, as well as decreased welfare. As an example: in the last four years, ASS has been causing problems in one BKD infected farm and mortalities continued despite antibiotic treatment. The reason is probably the underlying BKD infection, facilitating the ASS infection and itself being accelerated by the concurrent ASS infection. Control of BKD is expected to be improved by modified sampling and improved methodology, from today's post mortem sampling to an *in vivo* method. Additional resources must be invested in risk-based analysis of individual aquaculture farms to get a more reliable assessment for health surveillance.

Table 33: Samples taken in the Swedish surveillance programmes for notifiable diseases in fish, crustaceans and molluscs. One case=one outbreak.

Disease	No. of sampled production sites	No. of infected production sites	No. of tested individuals	No. of tested pools	No. of infected individuals/pools
Fish					
VHS	64	0	-	431	-/0
IHN	64	0	-	431	-/0
IPN	64	1	-	431	-/1
SVC	2 ^A	0	7 ^A	6	0/0
KHV	2 ^A	0	7	5 ^B	0/0
BKD	73 ^C	7 ^C	3363	-	32/-
Crustaceans					
<i>Aphanomyces astaci</i>	15 ^D	4 ^D	30	-	10/-
WSSv	0	-	-	-	-/-
Molluscs					
<i>Bonamia ostreae</i> ^E	4	0	150	0	0/-
<i>Marteilia refringens</i> ^E	4	0	150	0	0/-

^A One import company. No aquaculture farms were tested.

^B Pools of 2 individuals.

^C one production site represents a wild arctic char.

^D A total of 15 locations were sampled, representing 10 separate waterways with wild crayfish and one laboratory with crayfish. Four waterways were positive.

^E This sampling was performed as part of a project within the European Sea and Fisheries Fund.

Abbreviations:

VHS	Viral haemorrhagic septicemia
IHN	Infectious Haematopoietic Necrosis
IPN	Infectious pancreatic necrosis
SVC	Spring viraemia of carp
KHV	Koi herpesvirus
BKD	Bacterial Kidney Disease